


THIOLUX[®] Jet



Dry Flowable Micronized Sulfur

<i>Active Ingredient:</i>	
Sulfur	80.0%
<hr/>	
<i>Other Ingredients:</i>	20.0%
<hr/>	
<i>Total:</i>	100.0%
<hr/>	
<i>Guaranteed Analysis:</i>	
Sulfur(s)	80.0%
<i>Derived from elemental sulfur</i>	

**KEEP OUT OF REACH OF CHILDREN.
CAUTION**

OMRI[™]
L i s t e d

See additional precautionary statements and directions
for use printed on bag.

EPA Reg. No. 100-1138 EPA Est. 70992-FRA-002

Made in France

PRODUCT ID.

19532

SCP 1138A-L2B 1203

L110264 USA/12G

50 pounds

Net Weight

syngenta

Thiolux® Jet

FIRST AID	
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER For 24 Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident) Call 1-800-888-8372	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wear long-sleeved shirt and long pants, socks, shoes and chemical resistant gloves made of waterproof material. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.

Personal Protective Equipment

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves made of waterproof material
- Shoes plus socks
- Protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

Environmental Hazards

For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment and when disposing of equipment washwaters.

Physical and Chemical Hazards

Sulfur dust suspended in air ignites easily. Keep away from heat, sparks, and flame. Do not smoke while applying this product.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and Buyer and User assume the risk of any such use. SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

Thiolux® Jet

In no event shall SYNGENTA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls over long-sleeved shirt and long pants
- Chemical resistant gloves made of waterproof material
- Socks and chemical resistant footwear

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND THE PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR PEST CONTROL, OR ILLEGAL RESIDUES.

Thiolux Jet is recommended for control of listed plant diseases, insects and spider mites on fruits, vegetables, and ornamentals when used as directed. Additionally, Thiolux Jet is a readily available source of sulfur, a necessary plant nutrient.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Must be stored under dry conditions.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

Some crops are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use on any crop unless local use has proven sulfur to be safe in that locality. Sulfur may burn foliage or cause fruit injury when temperatures are excessively high. Do not apply at such times. Do not apply to or allow spray drift on sulfur sensitive plants such as apricots, cranberries, spinach. Apply only to dormant Anjou pears. Do not use within two (2) weeks of an oil treatment, except for dormant sprays on pears and apples. For citrus, do not apply within 21 days of an oil application. Follow all other label directions or precautions as indicated. When growing crops for processing, consult processor before applying sulfur.

Thiolux® Jet

Nutrient Use

Thiolux Jet applications will provide sulfur for the plant's nutrient requirements and should be considered in the total plant nutrient applications. Thiolux Jet can be used as a micronutrient supplement by either foliar or soil application. Thiolux Jet can be used on the following crops in addition to other crops listed on the label: alfalfa, almonds, avocados, apples, barley, all beans, broccoli, cabbage, cauliflower, carrots, citrus, corn, grapes, grass grown for seed, melons, nectarines, pears, peaches, peppers, plums, prunes, potatoes, peanuts, sorghum, soybeans, sugar beets, strawberries, tomatoes, turnips, walnuts, watermelons, and wheat. Consult your State Agricultural Experimental Station or Extension Specialist for advice in selecting treatments to best fit local conditions.

APPLICATION PROCEDURES

Fill spray tank half full with water. Pour needed amount into spray tank under agitation. If necessary, add a few drops of oil to avoid foaming. Maintain agitation throughout spray operation.

Use lower recommended rates on following host and pest recommendation table when using concentrate equipment. Use higher rates when using dilute applications.

Dilute Application

Ground Application – Apply specified rate in 20-60 gals. of water per acre.

Orchard Application – Apply specified rate in 100-800 gals. of water per acre.

Concentrate Application

Ground Application – Apply specified rate in 5 gals. or more water per acre.

Orchard Application – Apply specified rate in 20-100 gals. of water per acre. Special concentrate equipment is necessary for these uses.

Aerial Application

Apply specified rate in 3-20 gals. of water per acre. Orchard rates should not be applied in less than 10 gals. of water per acre.

Chemigation

Thiolux Jet may be applied through sprinkler systems including center pivot, lateral move, end tow, side wheel roll, traveler, big gun, solid set, and hand move. Do not apply Thiolux Jet through any other type irrigation system. Do not apply Thiolux Jet through any irrigation system when wind speed favors drift beyond the area intended for treatment.

Crop injury or lack of effectiveness can result from uneven distribution of treated irrigation water. For center pivot systems, apply during one full revolution of the system. For other systems listed above (i.e., portable or solid set systems), apply during the final 15 minutes of the irrigation period. Do not operate irrigation system after application until deposit has thoroughly dried. If you have questions about system operation or calibration, contact State Extension Service specialists or chemigation equipment manufacturers.

THE FOLLOWING PRECAUTIONS MUST BE OBSERVED WHEN APPLYING PESTICIDES AND PLANT NUTRIENTS THROUGH IRRIGATION SYSTEMS:

- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. Public water system means any system for provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
- Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- The pesticide injection system must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

THIOLUX JET APPLICATION RATES BY CROP AND PEST

Field Crops, Orchard Crops, Vegetables

Crop	Pest	lbs./A
alfalfa, cotton	Atlantic mite, red spider mite	3-5
Apply as needed.		
almonds, cherries, nectarines, peaches, plums, prunes	brown rot, blossom blight, cherry leafspot, powdery mildew, rust, scab, leafspot, silver mite, flat mite, red spider mite	10-30
Apply at bloom or early petal fall. Repeat as necessary, usually 10-14 days after a period of wet weather. For brown rot of fruit, apply when fruit starts to ripen. For rust of peaches and prunes, apply anytime from late spring to fall in a preventative program.		
apples, pears	powdery mildew, scab, blister mite, red spider mite	10-30
Apply at pre-bloom and petal fall. Apply with cover sprays throughout the season. Dormant applications with oil may be made to apples and pears. Apply to Anjou pears only while dormant.		
asparagus	rust	10-30
Apply after cutting stops. Irrigate and cultivate before applying; repeat at 7 to 10-day intervals throughout the season.		
avocados	brown mite	10-30
Apply as needed.		
barley, corn, oats, rye, sorghum, soybean, wheat	suppression – red spider mite, two-spotted mite, grass banks mite, Pacific mite; control – Atlantic mite (small grain only) powdery mildew (soybean only)	6 7.5-12
Apply when disease first appears and repeat as necessary or with the regularly scheduled insect control program.		
beans, peas, potatoes, rutabagas, turnips	powdery mildew, rust, red spider mite	3-10
Begin applications in early leaf stage. Repeat application every 10-14 days with or without pesticides immediately after rainfall.		
broccoli, brussel sprouts, cabbage, cauliflower, collards, kale, peppers	powdery mildew, rust, red spider mite	3-5
Begin applications in early leaf stage. Repeat application every 10-14 days with or without pesticides immediately after rainfall.		
blackberries, boysenberries, gooseberries, loganberries	powdery mildew	6-15
Begin application when new shoots are 6 inches long and before blossoms open or when disease first appears. Repeat at 7 to 10-day intervals or as necessary.		
blueberries, currants, dewberries, huckleberries	powdery mildew	5-10
Begin application when new shoots are 6 inches long and before blossoms open or when disease first appears. Repeat at 7 to 10-day intervals or as necessary.		
carrots, garlic, onions	Petrobia mite, powdery mildew	3-10
Apply when the disease first appears and repeat as needed.		
citrus	flat mite, aardvark mite, silver mite, rust mite, bud mite, red spider mite, six spotted mite, thrips, yuma spider mite	8-20
Apply in any spray NOT CONTAINING OIL or 21 days following oil treatment.		
cucurbits – cucumbers melons squash	powdery mildew powdery mildew powdery mildew	4-6 6-15 4-10
Begin application when disease first appears. Repeat at 10 to 14-day intervals or as necessary throughout the growing season. Sulfur can injure plants, especially when temperatures reach 95°F. Do not use on sulfur sensitive varieties.		
grapes	powdery mildew, bud mite, blister mite, red spider mite	6
Begin application in late dormant stage or with cutworm spray in early shoot period. Repeat throughout season on 14-day schedule or with pesticides.		
mangoes	powdery mildew	10-30
Begin applications before bloom, repeat in-bloom, after fruit set, and 3 weeks later or as necessary.		
peanuts	rust, red spider mite, leafspot	3-5
Begin application in early leaf stage. Repeat applications as necessary, usually in 10-24 days.		
peppermint, spearmint	powdery mildew	4-6
Apply when mint is 5-6 inches tall or when disease appears. Repeat twice at 30-day intervals. Do not apply within 30 days of harvest.		

continued...

Thiolux® Jet

THIOLUX JET APPLICATION RATES BY CROP AND PEST (continued)

Field Crops, Orchard Crops, Vegetables

Crop	Pest	lbs./A
pistachio	citrus flat mite	10-30
Begin application when mites are first detected. Repeat applications as necessary.		
raspberries	powdery mildew	6-15
Apply weekly starting at first bloom color. Continue throughout fruit set; repeat as weather requires.		
strawberries	powdery mildew, red spider mite	5-10
Begin application in early leaf stage, repeat application with or without pesticides. For powdery mildew, begin at first signs of disease and repeat as necessary. Do not use on sulfur sensitive varieties.		
sugar beets	powdery mildew, red spider mite	10-20
Apply at 10 to 30-day intervals with or without other pesticides at first symptoms.		
table beets	powdery mildew, red spider mite	3-5
Apply at 10 to 30-day intervals with or without other pesticides at first symptoms.		
tomatoes	russet mite, powdery mildew	6-20
Apply at first symptoms. Thorough coverage is required.		
walnuts	Pacific mite, almond mite, two-spotted mite, red spider mite	10-20
Apply and repeat as necessary throughout season as necessary. Tank mix with chemical mite sprays to increase effectiveness. NOTE – Certain varieties of walnuts are sensitive to sulfur sprays under certain conditions. Do not apply unless the varieties are known to be tolerant of sulfur.		

Ornamentals, Grass Seed Production

Ornamentals – all (see below for host and pest)		(ground) all → 3-10 (air) all → 3-20
Apply as aqueous spray for complete plant coverage. Begin when disease first appears and repeat at 5 to 10-day intervals as necessary.		
asters, carnations, cosmos, crepe myrtle, dahlia, delphinium, gladiolus, ligustrum, sunflowers, sweet peas, violets, zinnias	powdery mildew	
calendula, cherry laurels, chrysanthemums, hollyhock, smilax	leafspot, powdery mildew	
clematis, columbine, English ivy, laurel, petunia, sage	leafspot	
Roses	black spot, powdery mildew	
grass grown for seed	rust	3-5
Apply in tank mix with Tilt® for enhanced control of rusts. Begin applications when rust pustules are evident and increasing in number from late spring through early summer. Do not exceed 12 lbs./A of Thiolux Jet per season. Read and follow all application rates and use precautions on companion fungicide label. Do not apply this tank mix through any irrigation system.		

Forestry Uses

Douglas Fir trees – For use in Oregon Only.	Swiss Needle Cast (caused by <i>Phaeocryptopus gaeumannii</i>)	30-60
Thiolux Jet may provide suppression of Swiss needle cast caused by <i>Phaeocryptopus gaeumannii</i> . For best results, make the first application in the spring where there is 1-2" of new growth. Repeat as necessary every 10-14 days. Do not exceed a total of 3 applications per season. Use of a spreader-sticker is recommended to improve efficacy. Apply 30-60 lbs. of product per acre in sufficient water to obtain thorough, uniform coverage of the foliage. If making only one aerial pass over timber, use a minimum water volume of 15 gallons per acre. To improve coverage, apply 1/2 of the recommended rate per acre in a minimum of 10 gallons of water per acre one direction and repeat in the opposite direction using the remaining 1/2 of the recommended rate in a minimum of 10 gallons of water per acre. Do not exceed a total of 180 lbs. of Thiolux Jet per acre per year. Use the higher use rate under heavier disease pressure. Reapply the treatments the following year to protect the new foliage. Consult your local agricultural extension agent for correct timing of applications.		
Note: Aerial application only. For application only to trees intended for timber production. Do not apply to trees in seedling nurseries, or trees intended for Christmas tree, ornamental, or landscape usage.		

Oregon Only

Information regarding the contents and levels of metals in this product is available at the Oregon Department of Agriculture internet site:
<http://oda.state.or.us/fertilizer>

Washington State Only

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.agr.wa.gov/>.

Thiolux® Jet, Tilt® and the Syngenta logo are trademarks of a Syngenta Group Company

OMRI™ is a trademark of Organic Materials Review Institute

©2003 Syngenta

For non-emergency (e.g. additional product information) call
Syngenta Crop Protection at 1-800-334-9481.

Syngenta Crop Protection, Inc.
Greensboro, North Carolina 27409
www.syngenta-us.com

SCP 1138A-L2B 1203
L110264 USA/12G