

---

## SUPPLEMENTAL LABELING

---

**Syngenta Crop Protection, LLC**  
P. O. Box 18300  
Greensboro, North Carolina 27419-8300  
**SCP 1278A-S4 1210**

**Revus Top®**

**Fungicide**

Active Ingredients:	
Mandipropamid (CAS No. 374726-62-2).....	21.9%
Difenoconazole (CAS No.119446-68-3).....	21.9%
<hr/>	
Other Ingredients:	56.2%
<hr/>	
Total:	100.0%

Contains 2.08 pounds of mandipropamid active ingredient and 2.08 pounds of difenoconazole active ingredient per gallon

**KEEP OUT OF REACH OF CHILDREN.**

## **CAUTION**

**EPA Reg. 100-1278**

**All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using Revus Top as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and/or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

**syngenta**

## DIRECTIONS FOR USE

Crop	Target Disease	Use Rate fl oz Product per Acre	Remarks
<b>Grapes</b> (except Concord, Concord Seedless, and Thomcord. See Precaution under Remarks.)	Alternaria rot ( <i>A. alternata</i> )	7.0	For powdery mildew, begin at bud break and apply on a 10-21 day interval, making no more than 2 sequential applications before alternating to a fungicide with a different mode of action. For Phomopsis diseases, apply at bud break, before shoots are 0.5 inches in length, and then again when shoots are 5-6 inches in length.  For black rot, begin when shoot length is 1-3 inches and continue on a 10 day interval.  For all other diseases, begin applications prior to disease onset when conditions are conducive for disease. Apply Revus Top on a 10-14 day schedule making no more than 2 sequential applications before alternating to another fungicide with a different mode of action.  If disease pressure is high, use the shortest interval.  The addition of a spreading/penetrating type adjuvant such as a non-ionic surfactant or crop oil concentrate or blend is recommended when applying by ground or air.  <b>PRECAUTION: On <i>V. labrusca</i>, <i>V. labrusca</i> hybrids, and other non-vinifera hybrids where sensitivity is not known - the use of Revus Top by itself or in tank mixtures with materials that may increase uptake (adjuvants, foliar fertilizers) may result in leaf burning or other phytotoxic effects.</b>
	Powdery mildew ( <i>Uncinula necator</i> )		
	Rotbrenner ( <i>Pseudopezicula tracheiphila</i> )		
	Septoria leaf spot ( <i>S. ampelina</i> )		
	Phomopsis cane and leaf spot ( <i>P. viticola</i> )		
	Black rot ( <i>Guignardia bidwellii</i> )		
	Angular leaf spot ( <i>Mycosphaeraella angulata</i> )		
	Anthracnose ( <i>Elsinoe ampelina</i> )		
	Leaf blight ( <i>Pseudocercospora vitis</i> )		
Downy mildew ( <i>Plasmopara viticola</i> )			

**Application:** For best results, sufficient water volume must be used to provide thorough coverage. Revus Top can be applied by either ground or aerial application. A minimum of 15 gal/A for ground applications is recommended. For aerial applications a minimum of 10 gal/A of water is recommended.

**Specific Use Restrictions:**

- 1) Do not apply more than 28 fl oz/A of Revus Top per crop per season.
- 2) Do not apply more than 0.46 lb ai/A per season of difenoconazole containing products.
- 3) Do not apply more than 0.52 lb ai/A per season of mandipropamid containing products.
- 4) Do not apply within 14 days of harvest (14-day PHI).

Revus Top® and the Syngenta logo are trademarks of a Syngenta Group Company

©2011 Syngenta

REV TOP 1278A-S4 1210 – sw – 3/18/11